

## Acetaminophen Suppositories (rectal)

FeverAll<sup>®</sup>

- You may give your child acetaminophen if your child is uncomfortable with a fever.
- Acetaminophen should **NOT** be used in children under 2 months
- Use the acetaminophen dosing chart below. Use your child's **weight** to select the dose.
- You may repeat the dose of acetaminophen every 4-6 hours.

Weight	Age	Infant's 80 mg	Children's 120 mg
<12 lbs	< 6 months		
12-17 lbs	6-11 months	1 suppository	–
18-23 lbs	12-23 months	–	1 suppository
24-35 lbs	3-6 years	–	1½ suppositories

### FeverAll Suppository Directions

- When practical, cleanse area around rectum with mild soap and warm water and rinse thoroughly.
- Gently dry by patting or blotting with toilet tissue or a soft cloth before using FeverAll.
- Remove wrapper before inserting suppository and hold suppository upright and carefully separate wrapper at pointed end to open.
- Carefully peel wrapper down both sides evenly, exposing suppository.
- Avoid excess handling, as suppository is designed to melt at body temperature.
- Suppositories should not need to be refrigerated if stored below 80° F (27° C).
- Place child flat on back or on one side with anal opening exposed.
- Insert suppository gently, well into the rectum. Use fingertip to complete insertion.